

# Leaky Gut's Contribution to Inefficient Nutrient Utilization

S.K. Kvidera<sup>1</sup>, E.A. Horst<sup>1</sup>, M. Al-Qaisi<sup>1</sup>, M.J. Dickson<sup>1</sup>, R.P. Rhoads<sup>2</sup>, and L.H. Baumgard<sup>1</sup>

<sup>1</sup>Iowa State University Department of Animal Science

<sup>2</sup>Virginia Tech Department of Animal Science

Corresponding author: baumgard@iastate.edu

## INTRODUCTION

There are a variety of situations in an animal's life when nutrient utilization is reprioritized from productive towards agriculturally unproductive purposes. Two well-known examples that markedly reduce production are heat stress and ketosis. Decreased feed intake, experienced during both diseases, is unable to fully explain decreases in productivity. Additionally, both diseases are characterized by negative energy balance, body weight loss, inflammation, and hepatic steatosis. While the metabolism of ketosis and heat stress have been thoroughly studied for the last 40 years, the initial insult in the cascade of events ultimately reducing productivity in both heat-stressed and ketotic cows has not been identified. To that end, we have generated preliminary data strongly implicating a metabolic disruptor, endotoxin, as the etiological culprit in each case.

### *Heat Stress*

Heat stress negatively impacts a variety of production parameters and is a significant financial burden (~\$900 million/year for dairy in the U.S. alone; St. Pierre et al., 2003). Heat-stress affects productivity indirectly by reducing feed intake; however, direct mechanisms also contribute as we have shown reduced feed intake only explains approximately 35-50% of the decreased milk yield during heat stress (Rhoads et al., 2009; Wheelock et al., 2010; Baumgard et al., 2011). Direct mechanisms contributing to heat stress milk yield losses involve an altered endocrine profile, including reciprocal changes in circulating anabolic and catabolic hormones (Bernabucci et al., 2010; Baumgard and Rhoads, 2012). Such changes are characterized by increased circulating insulin concentration, lack of adipose tissue lipid mobilization, and reduced adipocyte responsiveness to lipolytic stimuli. Hepatic and skeletal muscle cellular bioenergetics also exhibit clear differences in carbohydrate production and use, respectively, due to heat stress. Thus, the heat stress response markedly alters post-absorptive carbohydrate, lipid, and protein metabolism through coordinated changes in fuel supply and utilization across tissues in a manner distinct from commonly recognizable changes that occur in

animals on a reduced plane of nutrition (Baumgard and Rhoads, 2013). The result of HS is underachievement of an animal's full genetic potential.

Ketosis

The periparturient period is associated with substantial metabolic changes involving normal homeostatic adaptations to support milk production. Unfortunately, a disproportionate amount of herd culling occurs before cows reach 60 days in milk (Godden, 2003). Ketosis is defined as an excess of circulating ketone bodies and is characterized by decreases in feed intake, milk production, and increased risk of developing other transition period diseases (Chapinal et al., 2012). Epidemiological data indicate about 20% of transitioning dairy cows clinically experience ketosis (BHBA > 3.0 mM; Gillund et al., 2001) while the incidence of subclinical ketosis (>1.2 mM BHBA) is thought to be much higher (> 40%; McArt et al., 2012). Ketosis is a costly disorder (estimated at ~\$300 per case; McArt et al., 2015) and thus it represents a major hurdle to farm profitability. Traditionally, ketosis is thought to result from excessive adipose tissue mobilization (Baird, 1982; Grummer, 1993; Drackley, 1999) which in turn contributes to fatty liver (hepatic steatosis) and excessive ketone body synthesis (Grummer, 1993).

## HEAT STRESS ETIOLOGY

Mechanisms responsible for altered nutrient partitioning during HS are not clear; however, they might be mediated by HS effects on gastrointestinal health and function as we and others have demonstrated HS compromised intestinal barrier function (Lambert et al., 2002; Dokladny et al., 2006; Pearce et al., 2013; Sanz-Fernandez et al., 2014). During HS, blood flow is diverted from the viscera to the periphery in an attempt to dissipate heat leading to intestinal hypoxia (Hall et al., 1999). Enterocytes are particularly sensitive to hypoxia and nutrient restriction (Rollwagen et al., 2006), resulting in ATP depletion and increased oxidative and nitrosative stress (Hall et al., 2001). This contributes to tight junction dysfunction and gross morphological changes that ultimately reduce intestinal barrier function (Lambert et al., 2002; Pearce et al., 2013). As a result, HS increases the passage of lu-

minal content into portal and systemic blood (Hall et al., 2001; Pearce et al., 2013). Endotoxin, otherwise referred to as lipopolysaccharide (LPS), is a glycolipid embedded in the outer membrane of Gram-negative bacteria, which are abundant and prolific in luminal content, and is a well-characterized potent immune stimulator in multiple species (Berczi et al., 1966; Giri et al., 1990; Tough et al., 1997). Activation of the immune system occurs when LPS binding protein (LBP) initially binds LPS and together with CD14 and TLR4 delivers LPS for removal and detoxification, thus LBP is frequently used as a biomarker for LPS infiltration (Cecilian et al., 2012). For a detailed description of how livestock and other species detoxify LPS see our recent review (Mani et al., 2012). Endotoxin infiltration during HS into the bloodstream which was first observed by Graber et al. (1971), is common among heat stroke patients (Leon, 2007) and is thought to play a central role in heat stroke pathophysiology as survival increases when intestinal bacterial load is reduced or when plasma LPS is neutralized (Bynum et al., 1979; Gathiram et al., 1987). It is remarkable how animals suffering from heat stroke or severe endotoxemia share many physiological and metabolic similarities to HS, such as an increase in circulating insulin (Lim et al., 2007). Infusing LPS into the mammary gland increased (~2 fold) circulating insulin in lactating cows (Waldron et al., 2006). In addition, we intravenously infused LPS into growing calves and pigs and demonstrated >10 fold increase in circulating insulin (Rhoads et al., 2009; Stoakes et al., 2015c,d). Interestingly, increased insulin occurs prior to increased inflammation and the temporal pattern agrees with our previous in vivo data and a recent in vitro report (Bhat et al., 2014) suggesting LPS stimulates insulin secretion, either directly or via GLP-1 (Kahles et al., 2014). The possibility that LPS increases insulin secretion likely explains the hyperinsulinemia we have repeatedly reported in a variety of heat-stressed agriculture models (Baumgard and Rhoads, 2013). Again, the increase in insulin in both models is energetically difficult to explain as feed intake was severely depressed in both experiments.


## TRANSITION PERIOD INFLAMMATION

Recently, the concept that LPS impacts normal nutrient partitioning and potentially contributes to metabolic maladaptation to lactation has started to receive attention. Although LPS itself has not been the primary causative focus, general inflammation has been the topic of investigations. Increased inflammatory markers following parturition have been reported in cows (Ametaj et al., 2005; Bertoni et al., 2008; Humblet et al., 2006; Mullins et al., 2012). Presumably, the inflammatory state following calving disrupts normal nutrient partitioning and is detrimental to productivity (Lor et al., 2005; Bertoni et al.,

2008), and this assumption was recently reinforced when TNF $\alpha$  infusion decreased productivity (albeit without overt changes in metabolism; Yuan et al., 2013; Martel et al., 2014). Additionally, in late-lactation cows, injecting TNF $\alpha$  increased (>100%) liver TAG content without a change in circulating NEFA (Bradford et al., 2009). Our recent data demonstrates increased inflammatory markers in cows diagnosed with ketosis only and no other health disorders. In comparison with healthy controls, ketotic cows had increased circulating LPS prior to calving and post-partum acute phase proteins such as LPS-binding protein, serum amyloid A, and haptoglobin were also increased (Fig. 1; Abuajamieh et al., 2015). Endotoxin can originate from a variety of locations, and obvious sources in transitioning dairy cows include the uterus (metritis), mammary gland (mastitis) and the gastrointestinal tract (Mani et al., 2012). However, we believe intestinal permeability may be responsible for inflammation observed in the transition dairy cow. A transitioning dairy cow undergoes a post-calving diet shift from a mainly forage based to a high concentrate ration. This has the potential to induce rumen acidosis which can compromise the gastrointestinal tract barrier (Khafipour et al., 2009).

In order to further investigate the effects of intestinal permeability on production and inflammation, we intentionally induced intestinal permeability in mid-lactation dairy cows using a gamma secretase inhibitor (GSI), a compound that specifically inhibits crypt stem cell differentiation into enterocytes via disrupting Notch signaling (van Es et al., 2005). We anticipated feed intake of GSI administered cows would decrease, so we pair-fed controls in order to eliminate the confounding effect of feed intake. Treatment with GSI decreased feed intake and altered jejunum morphology consistently with characteristics of leaky gut (shortened crypt depth, decreased villus height, decreased villus height to crypt depth ratio). Circulating insulin and LBP were increased in GSI cows relative to controls. Interestingly in our GSI model, acute phase proteins serum amyloid A and haptoglobin increased for both treatments over time, indicating inflammation was occurring in pair-fed controls as well (Stoakes et al., 2014). This is not surprising, as pair-fed controls were receiving ~20% of their ad libitum intake and decreased feed intake has been shown to increase intestinal permeability in feed restricted rodents and humans (Rodriguez et al., 1996; Welsh et al., 1998) and we have also observed this in pigs (Pearce et al., 2013; Sanz-Fernandez et al., 2014). Recently, we confirmed the detrimental effects of feed restriction in mid-lactation cows by demonstrating a linear increase in circulating acute phase proteins and endotoxin with increasing severity of feed restriction. Furthermore, cows fed 40% of ad libitum intake had shortened ileum villous height

and crypt depth, indicating reduced intestinal health (Stoakes et al., 2015b). In summary, inflammation is present during the transition period and likely contributes to changes in whole-animal energetics.




**Figure 1.** Markers of inflammation in healthy (solid line) and ketotic (dashed line) transition cows.

## METABOLISM OF INFLAMMATION

LPS-induced inflammation has an energetic cost which redirects nutrients away from anabolic processes that support milk and muscle synthesis (see review by Johnson, 1997, 1998) and thus compromises productivity and efficiency. Interestingly, immune cells become more insulin sensitive and consume copious amounts of glucose upon activation in order to support rapid proliferation and biosynthetic processes (Calder et al., 2007; Palsson-McDermott and O'Neill, 2013). In contrast, inflammation induces an insulin resistant state in skeletal muscle and adipose tissue (Liang et al., 2013; Poggi et al., 2007). Recent data has also demonstrated a decrease in ketone oxidation during LPS infiltration (Suagee et al., 2011; Friesard et al., 2015) which we believe may partly explain increased ketone body concentrations during the transition period.

Endotoxin has previously been recognized to be involved with metabolic dysfunction. In humans, both obesity and high fat diets are linked to endotoxemia (Cani et al., 2007, Gregor and Hotamisligil, 2011). Furthermore, LPS is involved with the development of fatty liver (Ilan, 2012), and cytokines are linked to lipid accumulation and cholesterol retention (Ma et al., 2008; Clément et al., 2008). Experimentally-induced endotoxemia in dairy cattle has been linked to several metabolic and endocrine disturbances including decreased circulating glucose, termination of pregnancy, leukopenia, disruption of ruminal metabolism, and altered calcium homeostasis (Griel et al., 1975; Giri et al., 1990; Waldron et al., 2003; Jing et al., 2014). The aforementioned pathological conditions are likely mediated by LPS-induced inflammation and the subsequent changes in nutrient partitioning (Fig. 2) caused by immune system activation.



**Figure 2.** LPS induced alterations in glucose metabolism and insulin sensitivity.

### *Energetic Cost of Immune Activation*

An activated immune system requires a large amount of energy and the literature suggests that glucose homeostasis is markedly disrupted (Leininger et al., 2000) during an endotoxin challenge. Upon immune system activation, immune cells switch their metabolism from oxidative phosphorylation to aerobic glycolysis, causing them to become obligate glucose utilizers in a phenomenon known as the Warburg Effect (Vander Hiden et al., 2009). Our group recently employed a series of LPS-euglycemic clamps to quantify the energetic cost of an activated immune system. Using this model, we estimated approximately 1 kg of glucose is used by the immune system during a 12 hour period in lactating dairy cows. Interestingly, on a metabolic body weight basis the amount of glucose utilized by LPS-activated immune system in lactating cows, growing steers and growing pigs were 0.64, 1.0, and 1.1 g glucose/kg BW<sup>0.75</sup>/h, respectively; Stoakes et al., 2015a,c,d). Increased immune system glucose utilization occurs simultaneously with infection-induced decreased feed intake: this coupling of enhanced nutrient requirements with hypophagia obviously decrease the amount of nutrients available for the synthesis of valuable products (milk, meat, fetus, wool). We and others have now demonstrated that both heat-stressed and ketotic animals have increased circulating markers of endotoxin and inflammation. We believe that the circulating LPS in both maladies originates from the intestine and thus both likely have an activated immune system. This activated systemic immune response reprioritizes the hierarchy of glucose utilization and milk synthesis is consequently deemphasized.

### **CONCLUSION**

Ketosis and heat stress are two of the most economically important pathologies which severely jeopardize the competitiveness of animal agriculture. Heat stress and ketosis affect herds of all sizes and every dairy region in country. The biology of ketosis and heat stress has been studied for almost a half century, but the negative impacts of both are as severe today as they were 30 years ago. We suggest, based upon the literature and on our supporting evidence, that LPS is the common culprit etiological origin of both metabolic disorders. Taken together, our data and the literature suggest that LPS markedly alters nutrient partitioning and is a causative agent in metabolic disruption during heat stress and ketosis.

\*Parts of this manuscript were first published in the proceedings of the 2016 Southwest Nutrition Conference in Tempe AZ.

### **REFERENCES**

- Abujamieh, M., S. K. Stoakes, M. V. Sanz Fernandez, J. S. Johnson, J. T. Seibert, E. A. Nolan, S. M. Lei, H. B. Green, K. M. Schoenberg, W. E. Trout, and L. H. Baumgard. 2015. Characterizing the temporal pattern of leaky gut biomarkers in healthy and ketotic cows during the transition period. *J. Dairy Sci.* 98(E-Suppl. 2):876.
- Ametaj, B. N., B. J. Bradford, G. Bobe, R. A. Nafikov, Y. Lu, J. W. Young, and D. C. Beitz. 2005. Strong relationships between mediators of the acute phase response and fatty liver in dairy cows. *Can. J. Anim. Sci.* 85:165–175.



- Baird, G. D. 1982. Primary ketosis in the high-producing dairy cow: clinical and subclinical disorders, treatment, prevention, and outlook. *J. Dairy Sci.* 65:1-10.
- Baumgard, L. H. and R. P. Rhoads. 2013. Effects of heat stress on postabsorptive metabolism and energetics. *Annu. Rev. Anim. Biosci.* 1:311-337.
- Baumgard, L. H., J. B. Wheelock, S. R. Sanders, C. E. Moore, H. B. Green, M. R. Waldron, and R. P. Rhoads. 2011. Postabsorptive carbohydrate adaptations to heat stress and monensin supplementation in lactating Holstein cows. *J. Dairy Sci.* 94:5620-5633.
- Baumgard, L. H., and R. P. Rhoads. 2012. Ruminant production and metabolic responses to heat stress. *J. Anim. Sci.* 90:1855-1865.
- Berczi, I., L. Bertok, and T. Bereznai. 1966. Comparative studies on the toxicity of *Escherichia coli* lipopolysaccharide endotoxin in various animal species. *Can. J. of Microbiol.* 12:1070-1071.
- Bernabucci, U., N. Lacetera, L. H. Baumgard, R. P. Rhoads, B. Ronchi, and A. Nardone. 2010. Metabolic and hormonal acclimation to heat stress in domesticated ruminants. *Animal* 4(7):1167-1183.
- Bertoni, G., E. Trevisi, X. Han, and M. Bionaz. 2008. Effects of inflammatory conditions on liver activity in puerperium period and consequences for performance in dairy cows. *J. Dairy Sci.* 91:3300-3310.
- Bhat, U. G., V. Ilievski, T. G. Unterman, and K. Watanabe. 2014. *Porphyromonas gingivalis* lipopolysaccharide (LPS) upregulates insulin secretion from pancreatic beta cells line MIN6. *J. Periodontol.* 85:1629-1636.
- Bradford, B. J., L. K. Mamedova, J. E. Minton, J. S. Drouillard, and B. J. Johnson. 2009. Daily injection of tumor necrosis factor- $\alpha$  increases hepatic triglycerides and alters transcript abundance of metabolic genes in lactating dairy cattle. *J. Nutr.* 139:1451-1456.
- Bynum, G., J. Brown, D. Dubose, M. Marsili, I. Leav, T. G. Pistole, M. Hamlet, M. LeMaire, and B. Caleb. 1979. Increased survival in experimental dog heatstroke after reduction of gut flora. *Aviat. Space Environ. Med.* 50:816-819.
- Calder, P. C., G. Dimitriadis, and P. Newsholme. 2007. Glucose metabolism in lymphoid and inflammatory cells and tissues. *Curr. Opin. Clin. Nutr. Metab. Care* 10:531-540.
- Cani, P. D., J. Amar, M. A. Iglesias, M. Poggi, C. Knauf, D. Bastelica, A. M. Neyrinck, F. Fava, K. M. Tuohy, C. Chabo, A. Waget, E. Delmée, B. Cousin, T. Sulpice, B. Chamontin, J. Ferrières, J. F. Tanti, G. R. Gibson, L. Casteilla, N. M. Delzenne, M. C. Alessi, and R. Burcelin. 2007. Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* 56:1761-1772.
- Ceciliani, F., J. J. Ceron, P. D. Eckersall, and H. Sauerwein. 2012. Acute phase proteins in ruminants. *J. Proteomics* 75:4207-4231.
- Chapinal, N., S. J. Leblanc, M. E. Carson, K. E. Leslie, S. Godden, M. Capel, J. E. Santos, M. W. Overton, and T. F. Duffield. 2012. Herd-level association of serum metabolites in the transition period with disease, milk production, and early lactation reproductive performance. *J. Dairy Sci.* 95:5676-5682.
- Clément, S., C. Juge-Aubry, A. Sgroi, S. Conzelmann, V. Paziienza, B. Pittet-Cuenod, C. A. Meier, and F. Negro. 2008. Monocyte chemoattractant protein-1 secreted by adipose tissue induces direct lipid accumulation in hepatocytes. *Hepatology* 48:799-807.
- Dokladny, K., P. L. Moseley, and T. Y. Ma. 2006. Physiologically relevant increase in temperature causes an increase in intestinal epithelial tight junction permeability. *Am. J. Physiol. Gastrointest. Liver Physiol.* 290: G204-G212.
- Drackley, J. K. 1999. Biology of dairy cows during the transition period: the final frontier? *J. Dairy Sci.* 82: 2259-2273.
- Frisard, M. I., Y. Wu, R. P. McMillan, K. A. Voelker, K. A. Wahlberg, A. S. Anderson, N. Boutagy, K. Resendes, E. Ravussin, and M. W. Hulver. 2015. Low levels of lipopolysaccharide modulate mitochondrial oxygen consumption in skeletal muscle. *Metabolism* 64:416-427.
- Gathiram, P., M. T. Wells, J. G. Brock-Utne, and S. L. Gaffin. 1987. Antilipopolysaccharide improves survival in primates subjected to heat stroke. *Circ. Shock* 2:157-164.
- Gillund, P., O. Reksen, Y. T. Gröhn, and K. Karlberg. 2001. Body condition related to ketosis and reproductive performance in Norwegian dairy cows. *J. Dairy Sci.* 84:1390-1396.
- Giri, S. N., P. Emau, J. S. Cullor, G. H. Stabenfeldt, M. L. Bruss, R. H. Bondurant, and B. I. Osburn. 1990. Effects of endotoxin infusion on circulating levels of eicosanoids, progesterone, cortisol, glucose and lactic acid, and abortion in pregnant cows. *Vet. Microbiol.* 21:211-231.
- Godden, S. M., S. C. Stewart, J. F. Fetrow, P. Rapnicki, R. Cady, W. Weiland, H. Spencer, and S. W. Eicker. 2003. The relationship between herd rbST supplementation and other factors and risk for removal for cows in Minnesota Holstein dairy herds. Pages 55-64 in *Proc. Four-State Nutrition Conference. MidWest Plan. Service, LaCrosse, WI.*
- Graber, C. D., R. B. Reinhold, J. G. Breman, R. A. Harley, and G. R. Hennigar. 1971. Fatal heat stroke. Circulating endotoxin and gram-negative sepsis as complications. *JAMA* 216:1195-1196.
- Gregor, M. F. and G. S. Hotamisligil. 2011. Inflammatory mechanisms in obesity. *Annu. Rev. Immunol.* 29:415-445.
- Griel, L. C., A. Zarkower, and R. J. Eberhart. 1975. Clinical and clinico-pathological effects of *Escherichia coli* endotoxin in mature cattle. *Can. J. Comp. Med.* 39:1-6.