


# Biosecurity Checkpoints

## Practical tips for poultry farm biosecurity



### Terminal Disinfection

Empty shed cleaning followed by disinfection and fogging is a critical step for eliminating pathogens from shed.



### Regular Disinfection

In challenged farms disinfectant spray (triple salt or QUAT + Glutaraldehyde combination) should be done daily to reduce the spread of the disease.



### Pest Control Program

Implementation of pest control program for wild birds, rodents and insects should be followed strictly.



### Water Sanitization

Water sanitization should be done continuously for drinking water. Combination of acidifier and sanitizer can be used in the farms with high pH (more than 7).



### Restricted Entry of Visitors

Entry of essential personnel should only be allowed inside the farm.



### Minimize vehicle and equipment entry

Entry of vehicles and equipments should be minimized to farm. Washing and disinfecting the vehicle tyres and equipments should be done before entering the farm.



### Foot Bath

Feet should be dipped in the foot bath before entering and exiting the poultry shed. The foot bath should be changed twice in a day.



### Dead bird Disposal

Cremation or composting are the preferred methods for dead bird disposal. Proper cleaning and disinfection is recommended on the site of disposal.

## Keciddal™ T Plus

3 in 1 Farm & Hatchery  
Disinfectant Sanitizer

## Kem V 260®

Highly Effective Phenolic  
Disinfectant

## Trysil™

Highly Potent  
Broad Spectrum Virucidal

# KEMIN®

© Kemin Industries, Inc. and its group of companies 2021. All rights reserved. ®™ Trademarks of Kemin Industries, Inc., U.S.A.  
Certain statements, product labeling and claims may differ by geography or as required by government requirements.

